

Stern, D., Eisenhardt, P., Gorjian, V., Kochanek, C. S., Caldwell, N., Eisenstein, D., et al. (2005). Mid-Infrared Selection of Active Galaxies. *The Astrophysical Journal* , 631, 163-168.

Strasburger, D., Gorgian, V., Burke, T., Childs, L., Odden, C., Tambara, K., et al. (2015). Identification and Classification of Infrared Excess Sources in the Spitzer Enhanced Imaging Products (SEIP) Catalog. *AAS Meeting 225 Abstracts* (p. 1). Kissimmee, FL: American Astronomical Society.

Urry, C. M., & Padovani, P. (1995). Unified Schemes for Radio-Loud Active Galactic Nuclei. *Publications of the Astronomical Society of the Pacific* , 803-845.

Watson, D., Denney, K. D., Vestergaard, M., & Davis, T. M. (2011). A New Cosmological Distance Measure Using AGN. *The Astrophysical Journal Letters* , 740, 5pp.

Wolk, S. J., Gunther, M. H., Poppenhaeger, K., Cody, A. M., Rebull, L. M., Forbrich, J., et al. (2015). YSOVAR: Mid-infrared Variability Among YSOs in the Star Formation Region GGD12-15. *The Astronomical Journal* , 150, 23pp.

Zitrin, A., Labbe, I., Belli, S., Bouwens, R., Ellis, R. S., Roberts-Borsani, G., et al. (2015). Lyman α Emission from a Luminous $z = 8.6$ Galaxy: Implications for Galaxies as Tracers of Cosmic Reionization. *Astrophysical Journal Letters* , 810, 1-6.